1. A membrane artificial lung for performing gas exchange between blood and a gas via the membrane by flowing the blood in one side of the membrane and flowing oxygen or an oxygen-containing gas in the other side of the membrane,

wherein said membrane comprises a hollow fiber membrane, said hollow fiber membrane comprising poly-4-methylpentene-1 and having an oxygen permeation rate $Q(O_2)$ at 25°C of from 1, x: 10^{-6} to 3 x 10^{-3} (cm³(STP)/cm²/sec cmHg) and an ethanol flux of from 0.1 to 100 ml/min·m²,

wherein said membrane has, in the side of the blood flow, a surface comprising an ionic complex derived from:

quaternary alignatic alkylammonium salts; and heparin or a heparin derivative, and

wherein said quaternary alkylammonium salts comprise a quaternary aliphatic alkylammonium salt having from 22 to 26 carbon atoms in total and a quaternary aliphatic alkylammonium salt having from 37 to 40 carbon atoms in total.

2. The membrane artificial lung according to claim 1, wherein said quaternary alkylammonium salt comprises from 5 to 35% by weight of a quaternary aliphatic

alkylammonium salt having from 22 to 26 carbon atoms in total and from 65 to 95% by weight of a quaternary aliphatic alkylammonium salt having from 37 to 40 carbon atoms in total.

3. The membrane artificial lung according to claim 1, wherein said quaternary aliphatic alkylammonium salt comprise a dimethyldidodecylammonium salt and a dimethyldioctadecylammonium salt.

42

ABSTRACT OF THE DISCLOSURE

A membrane artificial lung for performing gas exchange between blood and a gas via the membrane by flowing the blood in one side of the membrane and flowing oxygen or an oxygen-containing gas in the other side of the membrane, wherein said membrane comprises a hollow fiber membrane, said hollow fiber membrane comprising poly-4-methylpentene-1 and having an oxygen permeation rate $Q(O_2)$ at 25°C of from 1 x 10⁻⁶ to 3 x 10⁻⁷ (cm³(STP)/cm² sec cmHg) and an ethanol flux of from 0.1 to 100 ml/min·m², wherein said membrane has, in the side of the blood flow, a surface comprising an ionic complex derived from: 'quaternary aliphatic' alkylammonium salts; and heparin or a heparin derivative, and wherein said quaternary alkylammonium salts comprise a quaternary aliphatic alkylammonium salt having from 22 to 26 carbon atoms in total and a quaternary aliphatic alkylammonium salt having from 37 to 40 carbon atoms in total.